

A WORD HOPF ALGEBRA BASED ON THE SELECTION/QUOTIENT PRINCIPLE

NGUYEN HOANG-NGHIA

LIPN - Institut Galilée - Université Paris 13, Sorbonne Paris Cité,
99, avenue Jean-Baptiste Clément, F-93430 Villetaneuse, France
email: `nguyen.hoang@lipn.univ-paris13.fr`

Abstract

This paper introduces a Hopf algebra structure (modelled after matroids) on the linear span of the set of packed word with selection-quotient coproduct. Using the notion of irreducible element, we prove that WMat is free as an algebra.

1 Introduction

In algebraic combinatorics, one often associates algebraic structures with various sets of combinatorial objects. Such structures are, for example, Hopf algebras on trees, graphs, tableaux, matroids, words, etc

A first type of combinatorial Hopf algebra structure is constructed using the selection/quotient principle. This means that the comultiplication is of the form:

$$\Delta(S) = \sum_{\substack{A \subseteq S \\ + \text{Conditions}}} S[A] \otimes S/A, \quad (1)$$

where $S[A]$ is a substructure of S and S/A is a quotient.

Examples of such Hopf algebras are the Connes-Kreimer Hopf algebra of Feynman graphs, underlying the combinatorics of perturbative renormalization in quantum field theory [1] or in non-commutative Moyal quantum field theory [8], [9] (the interested reader may also refer to [12] for a short review on these algebras). For the sake of completeness, let us also mention that similar Hopf algebraic structures have been proposed [7], [11] for quantum gravity spin-foam models.

Matroid theory has been introduced by Whitney in [13]. A structure of matroid Hopf algebra is defined in [12], where the product is given by the direct sum operation and the coproduct is given by the selection/quotient principle.

A second type of combinatorial Hopf algebra structure relies on the selection/complementation principle. This means that the comultiplication is of the form:

$$\Delta(S) = \sum_{\substack{A \subseteq S \\ + \text{Conditions}}} S[A] \otimes [S - A]. \quad (2)$$

Examples of such Hopf algebras are the Loday-Ronco Hopf algebra of planar binary trees [5] or the Hopf algebra on matrix quasi-symmetric functions **MQSym** [2].

It would thus be interesting to find Hopf algebraic structures on words with a comultiplication not of type (2), but of type (1). This is the issue we address in this paper.

In this article, we introduce a structure Hopf algebraic, that we call WMat, on the set of packed words with product given by the shifted concatenation and coproduct given by the selection/quotient principle, thus imitating matroid theory Hopf algebra. Each letter x_j can be seen as the infinite column vector with

Date: July 30, 2012

1 at j^{th} place and other entries zero. WMat has the zero column as a special element. In graph theory, it would correspond to the self-loop. We prove that the new Hopf algebra WMat is a subalgebra of a non commutative extension of **MQSym**. Using the notion of irreducible element, we prove that WMat is free as an algebra. We compute the Hilbert coefficients of this Hopf algebra and the Hilbert series. The coefficients of the Hilbert series are twice the corresponding Bell numbers.

Let us mention that we use the term word-matroid for these packed words in which we define this matroid-inspired Hopf algebra.

2 Algebra structure

2.1 Definitions

Let X be the $\{x_i\}_{i \geq 0}$ (the alphabet) and X^* be the set of words with letters in the alphabet X . A word w of length $n = |w|$ is a mapping $i \mapsto w[i]$ from $[1..|w|]$ to X . For a letter $x_i \in X$, the partial degree $|w|_{x_i}$ is the number of times the letter x_i occurs in the word w . One has:

$$|w|_{x_i} = \sum_{j=1}^{|w|} \delta_{w[j], x_i} . \quad (3)$$

For a word $w \in X^*$, one defines the alphabet $Alph(w)$ as the set of its letters, while $IAlph(w)$ the set of indices in $Alph(w)$.

$$Alph(w) = \{x_i \mid |w|_{x_i} \neq 0\} ; IAlph(w) = \{i \in \mathbf{N} \mid |w|_{x_i} \neq 0\} . \quad (4)$$

The upper bound $sup(w)$ is the supremum of $IAlph(w)$, i. e.

$$sup(w) = sup_{\mathbf{N}}(IAlph(w)) . \quad (5)$$

Note that $sup(1_{X^*}) = 0$.

Let us define the substitution operators. Let $w = x_{i_1} \dots x_{i_m}$ and $\phi : IAlph(w) \longrightarrow \mathbf{N}$, with $\phi(0) = 0$. One then has:

$$S_{\phi}(x_{i_1} \dots x_{i_m}) = x_{\phi(i_1)} \dots x_{\phi(i_m)} . \quad (6)$$

Let u, v be two words; one defines the shifted concatenation $*$ by

$$u * v = uT_{sup(u)}(v) , \quad (7)$$

where, for $t \in \mathbf{N}$, $T_t(w)$ denotes the image of w by S_{ϕ} for $\phi(n) = n + t$ if $n > 0$ and $\phi(0) = 0$ (in general, all letters can be reindexed except x_0).

Definition 2.1 Let $w \in X^*$ and $I = IAlph(w) \setminus \{0\}$ be the set of indices of w . Let $I = \{j_1, \dots, j_k\}$ with $j_1 < j_2 < \dots < j_k$ and define ϕ_0 as $\phi_0(j_m) = m$ and $\phi_0(0) = 0$. The pack of word $pack(w)$ is $S_{\phi_0}(w)$. A word $w \in X^*$ will be said packed iff $w = pack(w)$.

The operator $pack : X^* \longrightarrow X^*$ is idempotent ($pack \circ pack = pack$). It defines, by linear extension, a projector. Its image, $pack(X^*)$, is the set of packed words.

Definition 2.2 Let k be a field. One defines a vector space $\mathcal{H} = span_k(pack(X^*))$. One can endow this space with a product (on the words) given by

$$\begin{aligned} \mu : \mathcal{H} \otimes \mathcal{H} &\longrightarrow \mathcal{H}, \\ u \otimes v &\longmapsto u * v. \end{aligned}$$

Remark 1 The product above is similar to the shifted concatenation for permutations (see [2]). Moreover, let u, v be two words in X^* , then $sup(u * v) = sup(u) + sup(v)$. One then has:

$$T_{sup(u*v)} = T_{sup(u)+sup(v)} \dots \quad (8)$$

Proposition 2.3 $(\mathcal{H}, \mu, 1_{X^*})$ is an associative algebra with unit (AAU).

Proof.

Let u, v, w be three words in $\mathcal{H}$. One then has:

$$(u * v) * w = (uT_{\sup(u)}(v))T_{\sup(u*v)}(w) = u(T_{\sup(u)}(v)(T_{\sup(u)+\sup(v)}(w))) = u * (v * w). \quad (9)$$

Thus, $(\mathcal{H}, \mu)$ is associative. On the other hand, for all $u \in \text{pack}(X^*)$, one has:

$$u * 1_{X^*} = uT_{\sup(u)}(1_{X^*}) = u1_{X^*} = u,$$

$$1_{X^*} * u = (1_{X^*})T_{\sup(1_{X^*})}(u) = (1_{X^*})u = u.$$

Now remark that $\text{pack}(1_{X^*}) = 1_{X^*}$. This is clear from the fact that $1_{X^*} = 1_{\mathcal{H}}$.

One concludes that $(\mathcal{H}, \mu, 1_{X^*})$ is an AAU. $\square$

Remark 2 The product is non-commutative. This can be seen from the following example: $x_1 * x_1 x_1 \neq x_1 x_1 * x_1$.

Lemma 2.4 Let u, v be two words. Let $I \subset [1 \dots |u|]$ and $J \subset [|u| + 1 \dots |u| + |v|]$. One then has

$$\text{pack}(u * v[I + J]) = \text{pack}(u[I]) * \text{pack}(v[J']), \quad (10)$$

where J' is the set $\{i - |u|\}_{i \in J}$.

Proof.

By direct computation, one has:

$$\text{pack}(u * v[I + J]) = \text{pack}(uT_{\sup(u)}(v)[I + J]) = \text{pack}(u[I]T_{\sup(u)}(v)[J]). \quad (11)$$

One further has $\sup(u[I]) \leq \sup(u)$, and this, together with (11) leads to:

$$\text{pack}(u[I]T_{\sup(u)}(v)[J]) = \text{pack}(u[I]T_{\sup(u[I])}(v[J'])) = \text{pack}(u[I]) * \text{pack}(v[J']). \quad (12)$$

$\square$

In particular, when one takes $I = [1 \dots |u|]$ and $J = [|u| + 1 \dots |u| + |v|]$. The mapping pack

$$k < X > \xrightarrow{\text{pack}} \mathcal{H}$$

is then a morphism AAU.

2.2 WMat as a subalgebra of an extension of MQSym

Definition 2.5 Let the matrix M have the block diagonal form

$$M = \begin{pmatrix} M_1 & 0 & \cdots & \cdots \\ 0 & M_2 & \cdots & \cdots \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & M_k \end{pmatrix},$$

with k maximal and M_i is not an empty matrix. One defines the "free basis" of **MQSym**

$$FB_M = \mathbf{MS}_{M_1} \mathbf{MS}_{M_2} \dots \mathbf{MS}_{M_k}.$$

Lemma 2.6 (i) For all packed matrices P, Q , one has:

$$FB_P FB_Q = FB \begin{pmatrix} P & 0 \\ 0 & Q \end{pmatrix}.$$

(ii) $(FB_P)_{P \text{ is packed matrix}}$ is a free base of **MQSym**.

Proof. The conclusions follows directly from Definition 2.5. □

Definition 2.7 Let k be a field. One defines two mappings:

1) Let $Y = \{y, y_0\}$ an alphabet. Let

$$\begin{aligned} \pi'_0 : X &\longrightarrow k\langle Y \rangle \\ x &\longmapsto \begin{cases} y & \text{if } x \in X_1 \\ y_0 & \text{otherwise} \end{cases} \end{aligned}$$

One then has $\pi'_1 = \pi_1 \circ \text{nat}$ in the following diagram:

$$\begin{array}{ccc} X & \xrightarrow{\pi'_1} & k\langle Y \rangle \\ & \searrow \text{nat} & \nearrow \pi_1 \\ & k\langle X \rangle & \end{array}$$

2) Let $X_1^* = X^* - \{x_0\}$ and

$$\begin{aligned} \text{proj} : \text{pack}(X^*) &\longrightarrow \text{pack}(X_1^*) \\ w &\longmapsto \tilde{w} \end{aligned}$$

One can define the mapping π'_2 from $\text{pack}(X_1^*)$ to **MQSym** as follows: a word $w \in \text{pack}(X_1^*)$ is associated to the finite matrix M_w of size $l(w) \times \text{sup}(w)$, where the j^{th} column is e_k if $w[j] = x_k$. This matrix is packed. One has: $\pi'_2(w) = FB_{M_w}$.

One then has $\pi_2 = \pi'_2 \circ \text{proj}$ in the following diagram:

$$\begin{array}{ccc} \text{pack}(X) & \xrightarrow{\pi_2} & \mathbf{MQSym} \\ & \searrow \text{proj} & \nearrow \pi'_2 \\ & \text{pack}(X_1) & \end{array}$$

One denotes the matrix $\pi_2(w) = FB_{\pi'_2(w)}$.

Let $\mathcal{A} = k\langle y, y_0 \rangle \otimes \mathbf{MQSym}$. This is an extension of **MQSym** that we use later on in this paper.

Lemma 2.8 A mapping

$$\begin{aligned} \mu : \text{pack}(X) &\longrightarrow \mathcal{A} \\ w &\longmapsto \pi_1(w) \otimes \pi_2(w) \end{aligned} \tag{13}$$

is an injective morphism.

Proof.

Let u and u' two words in WMat. Suppose that $\mu(u) = \mu(u')$. One has:

$$\begin{cases} \pi_1(u) = \pi_1(u') \\ \pi_2(u) = \pi_2(u') \end{cases}.$$

From the first equation, it follows that the sets of positions of x_0 in the two words are the same and the two words have the same length. From the second equation, it follows that $u[i] = u'[i]$, $\forall i$ such that $u[i] \neq x_0$ and $u'[i] \neq x_0$. One then has $u = u'$, i.e the mapping μ is injective.

Let u and v two packed words.

One has

$$\pi_1(u * v) = \pi_1(uT_{sup(u)}(v)) = \pi_1(u)\pi_1(T_{sup(u)}(v)) = \pi_1(u)\pi_1(v) ,$$

and


$$\pi_2(u * v) = \pi_2(uT_{sup(u)}(v)) = FB_{\pi'_2(uT_{sup(u)}(v))} = FB \begin{pmatrix} \pi'_2(u) & 0 \\ 0 & \pi'_2(v) \end{pmatrix} = FB_{\pi'_2(u)}FB_{\pi'_2(v)} = \pi_2(u)\pi_2(v) ..$$

One then conclude that μ is a morphism. $\square$

From this lemma, one can identify the algebra WMat with $Im(\mu)$. The elements of $Im(\mu)$ have the form: $u \otimes v$ satisfying $u|_y = \#\{\text{column of } v\}$. The algebra WMat can thus be shown to be a subalgebra of $\mathcal{A}$.

2.3 WMat is a free algebra

WMat is a vector space with basis $pack(X)$. In order to check that WMat is a free algebra, it is sufficient to show that $pack(X)$ is a free monoid on its letters. In the following diagram, a free monoid is a pair $(F(X), j_X)$ where $F(X)$ is a monoid, $j_X : X \rightarrow F(X)$ is a mapping such that $(\forall M \in Mon) (\forall f : X \rightarrow M) (\exists ! f_X \in Mor(F(X), M)) f = f_X \circ j_X$.



Here we will use an "internal" characterization of free monoids in terms of irreducible elements.

Definition 2.9 A packed word w in $pack(X)$ is called an irreducible word if and only if it can not be written under the form $w = u * v$ where u and v are two non trivial packed words.

Example 2.10 The word $x_1x_1x_1$ is an irreducible word. The word $x_1x_1x_2$ is a reducible word because it can be written as $x_1x_1x_2 = x_1x_1 * x_2$.

Lemma 2.11 Let w be a packed word in $pack(X)$, then w can be written uniquely as $w = v_1 * v_2 * \dots * v_n$ where $v_i, 1 \leq i \leq n$, are non-trivial irreducible words.

Proof. A word w can be separated as $w = u * v$ if and only if $sup(u) = inf(v) - 1$ or $sup(v) = 0$. The i^{th} position of word w is called an admissible cut if $sup(w[1 \dots i]) = inf(w[i + 1 \dots |w|]) - 1$ or $sup(w[i + 1 \dots |w|]) = 0$.

Because the length of word is finite, one can get $w = v_1 * v_2 * \dots * v_n$, with n maximal and v_i non trivial, $\forall 1 \leq i \leq n$.

One assumes that one word can be written in two ways

$$w = v_1 * v_2 * \dots * v_n \tag{14}$$

and

$$w = v'_1 * v'_2 * \dots * v'_m . \tag{15}$$

Denoting by k the first number such that $v_k \neq v'_k$, without loss of generality, one can suppose that $|v_k| < |v'_k|$. From equation (14), the k^{th} position is an admissible cut of w . From equation (15), the k^{th} position is not an admissible cut of w . One thus has a contradiction. One has $n = m$ and $v_i = v'_i$ for all $1 \leq i \leq n$. $\square$

One can thus conclude that $pack(X)$ is free as monoid on its letters.

3 Bialgebra structure

Let us give the definition of the coproduct and prove that the coassociativity property holds.

Definition 3.1 Let $A \subset X$, one defines $w/A = S_{\phi_A}(w)$ with $\phi_A(i) = \begin{cases} i & \text{if } x_i \notin A, \\ 0 & \text{if } x_i \in A \end{cases}$.

Let $w/u = w/Alph(u)$.

Definition 3.2 The coproduct of $\mathcal{H}$ is given by

$$\Delta(w) = \sum_{I+J=[1 \dots |w|]} \text{pack}(w[I]) \otimes \text{pack}(w[J]/w[I]), \forall w \in \mathcal{H}, \quad (16)$$

where this sum runs over all partitions of $[1 \dots |w|]$ divided into two blocks, $I \cup J = [1 \dots |w|]$ and $I \cap J = \emptyset$.

Example 3.3 One has:

$$\Delta'(x_1 x_1 x_1) = 3x_1 \otimes x_0^2 + 3x_1^2 \otimes x_0.$$

$$\Delta'(x_1 x_2) = 2x_1 \otimes x_1.$$

$$\Delta'(x_1 x_2 x_1) = x_1 \otimes x_1 x_0 + x_1 \otimes x_1^2 + x_1 \otimes x_0 x_1 + x_1 x_2 \otimes x_0 + x_1^2 \otimes x_1 + x_2 x_1 \otimes x_0.$$

Let us now prove the coassociativity.

Lemma 3.4 Let $w \in X^*$ be a word and let the subset $I \subset [1 \dots |u|]$ and the subset $I_1 \subset [1 \dots |I|]$. Let ϕ_I be the unique strictly increasing map from $IAlph(w[I])$ to $IAlph(\text{pack}(w[I]))$. One then has

$$\text{pack}(w[I])[I_1] = S_{\phi}(w[I_1']), \quad (17)$$

where I_1' is $\phi_I^{-1}(I_1)$.

Proof.

Let $I = [i_1, \dots, i_n]$, and $I_1 = [j_1, \dots, j_k] \subseteq [1 \dots |I|]$. Then $I_1' = [j_1', \dots, j_k'] = [\phi^{-1}(j_1), \dots, \phi^{-1}(j_k)]$. One has:

$$\text{pack}(w[I])[I_1] = w_{j_1} \dots w_{j_k}. \quad (18)$$

On the other hand, one also has:

$$S_{\phi_I}(w[I_1']) = w_{\phi_I(j_1')} \dots w_{\phi_I(j_k')} = w_{\phi_I(\phi_I^{-1}(j_1))} \dots w_{\phi_I(\phi_I^{-1}(j_k))} = w_{j_1} \dots w_{j_k}. \quad (19)$$

Using (18) and (19) leads to (17). $\square$

Lemma 3.5 Let $w \in X^*$ be a word and ϕ be a strictly increasing map from $IAlph(w)$ to $\mathbb{N}$. One then has:

$$1) \quad \text{pack}(S_{\phi}(w)) = \text{pack}(w). \quad (20)$$

$$2) \quad S_{\phi}(w_1 / w_2) = S_{\phi}(w_1) / S_{\phi}(w_2). \quad (21)$$

Proof.

1) One has

$$\text{pack}(S_{\phi}(w)) = S_{\phi_0}(S_{\phi}(w)) = S_{\phi_0 \circ \phi}(w),$$

where ϕ_0 is the packing map which is given in Definition 2.1.

Note that both ϕ and ϕ_0 are strictly increasing maps.

Denote $I = IAlph(w) = \{j_1, j_2, \dots, j_k\}$, $j_1 < j_2 < \dots < j_k$, the image set $\phi(I) = \{j'_i, j'_i = \phi(j_i), i = 1 \dots k\}$ one has $j'_1 < j'_2 < \dots < j'_k$. From the definition of ϕ_0 , one has: $\phi_0(j'_i) = i = \phi_0(j_i)$. This leads to:

$$S_{\phi_0 \circ \phi}(w) = S_{\phi_0}(w) = pack(w). \quad (22)$$

2) Let $I_2 = Alph(w_2)$ and $I'_2 = Alph(S_\phi(w_1))$.

Let us rewrite the two sides of equation (21), the left hand side (LHS) and the right hand side (RHS):

$$LHS = S_\phi(S_{\phi_{I_2}}(w_1)) = S_{\phi \circ \phi_{I_2}}(w_1), \quad (23)$$

$$RHS = S_{\phi_{I'_2}}(S_\phi(w_1)) = S_{\phi_{I'_2} \circ \phi}(w_1) \dots \quad (24)$$

With $x_i \in Alph(w_1)$, one has two cases:

1. If $x_i \in I_2$, then $\phi_{I_2}(i) = 0$ and $\phi \circ \phi_{I_2}(i) = 0$ because ϕ is a strictly increasing map.

On the other hand, $\phi(i) \in I'_2$ and this implies $\phi_{I'_2} \circ \phi(i) = \phi_{I'_2}(\phi(i)) = 0$.

2. If $x_i \notin I_2$, then $\phi_{I_2}(i) = i$ and $\phi \circ \phi_{I_2}(i) = \phi(i)$.

On the other hand, because ϕ is a strictly increasing map, then $\phi(i) \notin I'_2$, and $\phi_{I'_2} \circ \phi(i) = \phi(i)$.

One thus has $\phi \circ \phi_{I_2}(i) = \phi_{I'_2} \circ \phi(i)$. Using this result and the two equations above (23) and (24), we conclude the proof. $\square$

Lemma 3.6 *Let w be a word in $\mathcal{H}$, and I, J, K be three disjoint subsets of $\{1 \dots |w|\}$. One then has:*

$$\frac{w[K]/w[I]}{w[J]/w[I]} = w[K] / w[I+J]. \quad (25)$$

Proof.

Using Lemma 3.5, one has:

$$\frac{w[K]/w[I]}{w[J]/w[I]} = S_{\phi_I}(w[K]) / S_{\phi_I}(w[J]) = S_{\phi_I}(w[K]/w[J]) = S_{\phi_I}(S_{\phi_J}(w[K])) = S_{\phi_I \circ \phi_J}(w[K]) = w[K] / w[I+J]. \quad (26)$$

$\square$

Proposition 3.7 *The vector space $\mathcal{H}$ endowed with the coproduct (16) is coassociative. Moreover, let the counit $\epsilon \in \mathcal{H}^*$ such that:*

$$\epsilon(w) = \begin{cases} 1 & \text{if } w = 1_{X^*}, \\ 0 & \text{otherwise.} \end{cases}$$

Then $(\mathcal{H}, \Delta, \epsilon)$ is a coassociative coalgebra with counit (c-AAU).

Proof.

Let us first prove the coassociativity of the coproduct (16), namely

$$(\Delta \otimes Id) \circ \Delta(w) = (Id \otimes \Delta) \circ \Delta(w). \quad (27)$$

The LHS of the coassociativity condition (27) can be written:

$$\begin{aligned} (\Delta \otimes Id) \circ \Delta(w) &= \sum_{I+J=[1 \dots |w|]} \left(\sum_{I_1+I_2=[1 \dots |I|]} pack(pack(w[I])[I_1]) \otimes pack(pack(w[I][I_2])/pack(w[I][I_1])) \right) \\ &\otimes pack(w[J]/w[I]) = \sum_{I+J=[1 \dots |w|]} \left(\sum_{I'_1+I'_2=I} pack(S_\phi(u[I'_1])) \otimes pack(S_\phi(u[I'_2])/S_\phi(u[I'_1])) \right) \otimes pack(u[J]/u[I]) \\ &= \sum_{I'_1+I'_2+J=[1 \dots |w|]} pack(u[I'_1]) \otimes pack(u[I'_2]/u[I'_1]) \otimes pack(u[J]/u[I_1+I_2]). \end{aligned} \quad (28)$$

The RHS of the coassociativity condition (27) can be written as:

$$\begin{aligned}
(Id \otimes \Delta) \circ \Delta(w) &= \sum_{I+J=[1\dots|w|]} \text{pack}(u[I]) \otimes \left(\sum_{J_1+J_2=[1\dots|J|]} \text{pack}(\text{pack}(w^{[J]}/w[I]))[J_1] \right) \\
&\otimes \text{pack}(\text{pack}(\text{pack}(w^{[J]}/w[I])[J_2]/\text{pack}(w^{[J]}/w[I])[J_1])) = \sum_{I+J=[1\dots|w|]} \text{pack}(u[I]) \otimes \left(\sum_{J'_1+J'_2=J} \text{pack}(u^{[J'_1]}/u[I]) \right) \\
&\otimes \text{pack}(S_\phi(u^{[J'_2]}/u[I]/u^{[J'_1]}/u[I])) = \sum_{I+J=[1\dots|w|]} \text{pack}(u[I]) \otimes \left(\sum_{J'_1+J'_2=J} \text{pack}(u^{[J'_1]}/u[I]) \right) \\
&\otimes \text{pack}(u^{[J'_2]}/u_{[I+J'_1]}) = \sum_{I+J'_1+J'_2=[1\dots|w|]} \text{pack}(u[I]) \otimes \text{pack}(u^{[J'_1]}/u[I]) \otimes \text{pack}(u^{[J'_2]}/u_{[I+J'_1]}) \dots \quad (29)
\end{aligned}$$

Using the two equation (28) and (29), one can conclude that the coproduct (16) is coassociative. Let us now prove the following

$$(\epsilon \otimes Id) \circ \Delta(w) = (Id \otimes \epsilon) \circ \Delta(w), \quad (30)$$

for all word $w \in \mathcal{H}$.

Let us rewrite the LHS and RHS of the equation (30).

$$\begin{aligned}
LHS &= (\epsilon \otimes Id) \left(\sum_{I+J=[1\dots|w|]} \text{pack}(w[I]) \otimes \text{pack}(w[J]/w[I]) \right) \\
&= \sum_{I+J=[1\dots|w|]} \epsilon(\text{pack}(w[I])) \otimes \text{pack}(w[J]/w[I]) = 1_{X^*} \otimes \text{pack}(w) = \text{pack}(w) \dots \quad (31)
\end{aligned}$$

$$\begin{aligned}
RHS &= (Id \otimes \epsilon) \left(\sum_{I+J=[1\dots|w|]} \text{pack}(w[I]) \otimes \text{pack}(w[J]/w[I]) \right) \\
&= \sum_{I+J=[1\dots|w|]} \text{pack}(w[I]) \otimes \epsilon(\text{pack}(w[J]/w[I])) = \text{pack}(w) \otimes 1_{X^*} = \text{pack}(w) \dots \quad (32)
\end{aligned}$$

One thus concludes that $(\mathcal{H}, \Delta, \epsilon)$ is a coalgebra. $\square$

Remark 3 This coalgebra is not cocommutative. Let $u = x_1 x_2 x_2$. One then has:

$$\begin{aligned}
\tau \circ \Delta'(u) &= \tau(x_1 \otimes x_1 x_1 + 2x_1 \otimes x_1 x_0 + 2x_1 x_2 \otimes x_0 + x_1 x_1 \otimes x_1) \\
&= x_1 x_1 \otimes x_1 + 2x_1 x_0 \otimes x_1 + 2x_0 \otimes x_1 x_2 + x_1 \otimes x_1 x_1 \neq \Delta'(u).
\end{aligned}$$

Lemma 3.8 Let u, v be two words. Let $I_1 + J_1 = [1\dots|u|]$ and $I_2 + J_2 = [|u| + 1\dots|u| + |v|]$. One then has

$$\text{pack}(u^{*v[J_1+J_2]}/u^{*v[I_1+I_2]}) = \text{pack}(u^{[J_1]}/u_{[I_1]}) * \text{pack}(v^{[J'_2]}/v_{[I'_2]}), \quad (33)$$

where I'_2 is the set $\{k - |u|, k \in I_2\}$ and J'_2 is the set $\{k - |u|, k \in J_2\}$.

Proof.

One has:

$$\begin{aligned}
&\text{pack}(u^{*v[J_1+J_2]}/u^{*v[I_1+I_2]}) = \text{pack}(S_{\phi_{I_1+I_2}}(u * v[J_1 + J_2])) = \text{pack}(S_{\phi_{I_1+I_2}}(u T_{\text{sup}(u)}(v)[J_1 + J_2])) \\
&= \text{pack}(S_{\phi_{I_1} + \phi_{I_2}}(u[J_1] T_{\text{sup}(u)}(v)[J_2])) = \text{pack}(S_{\phi_{I_1}} S_{\phi_{I_2}}(u[J_1] T_{\text{sup}(u)}(v[J'_2]))) \\
&= \text{pack}(S_{\phi_{I_1}}(u[J_1]) S_{\phi_{I_2}}(T_{\text{sup}(u)}(v[J'_2]))) = \text{pack}(u^{[J_1]}/u_{[I_1]} T_{\text{sup}(u^{[J_1]}/u_{[I_1]})} \text{pack}(S_{\phi_{I'_2}}(v[J'_2]))) \\
&= \text{pack}((u^{[J_1]}/u_{[I_1]}) * \text{pack}(u^{[J'_2]}/u_{[I'_2]})) \dots \quad (34)
\end{aligned}$$

$\square$

Proposition 3.9 *Let u, v be two words in $\mathcal{H}$. One has:*

$$\Delta(u * v) = \Delta(u) *^{\otimes 2} \Delta(v) \quad (35)$$

Proof.

One has:

$$\begin{aligned} \Delta(u * v) &= \sum_{\substack{I_1 + I_2 = I \\ J_1 + J_2 = J \\ I_1, J_1 \subset [1 \dots |u|] \\ I_2, J_2 \subset [|u| + 1 \dots |u| + |v|]}} (\text{pack}(u * v[I_1 + I_2])) \otimes (\text{pack}^{(u * v[J_1 + J_2]) / (u * v[I_1 + I_2])}) \\ &= \sum_{\substack{I_1 + J_1 = [1 \dots |u|] \\ I'_2 + J'_2 = [1 \dots |v|]}} \left(\text{pack}(u[I_1]) \otimes \text{pack}(u^{[J_1]} / u[I_1]) \right) * \left(\text{pack}(v[I_2]) \otimes \text{pack}(u^{[J'_2]} / u[I'_2]) \right) \\ &= \left(\sum_{I_1 + J_1 = [1 \dots |u|]} \text{pack}(u[I_1]) \otimes \text{pack}(u^{[J_1]} / u[I_1]) \right) * \left(\sum_{I'_2 + J'_2 = [1 \dots |v|]} \text{pack}(v[I_2]) \otimes \text{pack}(u^{[J'_2]} / u[I'_2]) \right) \\ &= \Delta(u) *^{\otimes 2} \Delta(v) \quad (36) \end{aligned}$$

□

Since $\mathcal{H}$ is graded by the word's length, one has the following theorem:

Theorem 3.10 *$(\mathcal{H}, \Delta)$ is a Hopf algebra.*

Proof.

The proof follows from the above results. □

As already announced above, we call this Hopf algebra WMat (word-matroid).

The antipode is given by the formula:

$$S(w) = -w - \sum_{I+J=[1 \dots |w|], I, J \neq \emptyset} S(\text{pack}(w[I])) \text{pack}(w^{[J]} / w[I]) \quad (37)$$

4 Primitive elements of WMat

Lemma 4.1 *Prim(WMat) is a subalgebra of WMat, graded by the word's length:*

$$\text{Prim(WMat)}_n = \text{Prim(WMat)} \cap \text{WMat}_n \quad (38)$$

Proof. Let $p \in \text{Prim(WMat)}$. Since WMat is graded, one can write

$$p = \sum_{i \geq 1} p_i \quad (39)$$

where $|p_i| = i$.

One has:

$$\begin{aligned} \Delta(p) &= p \otimes 1 + 1 \otimes p = \left(\sum_{i \geq 1} p_i \right) \otimes 1 + 1 \otimes \left(\sum_{i \geq 1} p_i \right) = \sum_{i \geq 1} (p_i \otimes 1) + \sum_{i \geq 1} (1 \otimes p_i) \\ &= \sum_{i \geq 1} (p_i \otimes 1 + 1 \otimes p_i), \quad p \in \text{Prim(WMat)} \quad (40) \end{aligned}$$

On the other hand, one has:

$$\Delta(p) = \sum_{i \geq 1} \Delta(p_i) \quad (41)$$

From the two equations above, one has:

$$\Delta(p_i) = p_i \otimes 1 + 1 \otimes p_i . \quad (42)$$

Finally, one has $p_i \in \text{Prim}(\text{WMat})$ and this leads to the conclusion. $\square$

Let $\text{Prim}(\text{WMat})_n$ be the subalgebra generated by primitive elements with length n . One can see an element P of this subalgebra as a polynomial of packed words with length n :

$$P = \sum_{w \in \text{WMat}_n} \langle P|w \rangle w . \quad (43)$$

One has the following standard result:

Lemma 4.2 *Let u, v two primitive elements. The element $[u, v] = u * v - v * u$ is also primitive.*

The words x_0 and x_1 are primitive elements. From Lemma 4.2, the element $p_1 = [x_0, x_1] = x_0x_1 - x_1x_0$ is also a primitive element.

One can then define the general element in $\text{Prim}(\text{WMat})_{n+1}$ by the following recursion: $p_n = [x_0, p_n]$, for all $n \geq 1$.

Lemma 4.3 *One has: $p_n \neq 0, \forall n \geq 1$.*

Proof. One can prove by recursion the explicit formula for the element p_n . Let

$$P(n) : \quad p_n = \sum_{k=0}^n (-1)^k \binom{n}{k} x_0^{n-k} x_1 x_0^k , \text{ for all } n \geq 2 .$$

Indeed, with $n = 2$, one has:

$$p_2 = [x_0, p_1] = [x_0, x_0x_1 - x_1x_0] = x_0^2x_1 - 2x_0x_1x_0 + x_1x_0^2.$$

Suppose that $P(n)$ holds, one can check $P(n+1)$:

$$\begin{aligned} p_{n+1} &= [x_0, p_n] = [x_0, \sum_{k=0}^n (-1)^k \binom{n}{k} x_0^{n-k} x_1 x_0^k] = \sum_{k=0}^n (-1)^k \binom{n}{k} x_0^{n-k+1} x_1 x_0^k - \sum_{k=0}^n (-1)^k \binom{n}{k} x_0^{n-k} x_1 x_0^{k+1} \\ x_1 x_0^{k+1} &= x_0^{n+1} x_1 + \sum_{k=1}^n (-1)^k \left(\binom{n}{k} + \binom{n}{k-1} \right) x_0^{n-k+1} x_1 x_0^k + (-1)^{n+1} x_1 x_0^{n+1} \\ &= \sum_{k=0}^{n+1} (-1)^k \binom{n+1}{k} x_0^{n+1-k} x_1 x_0^k . \end{aligned}$$

This completes the proof. $\square$

From Lemma 4.2, it follows that p_n is primitive for all $n \geq 1$. One then has $\dim(\text{Prim}(\text{WMat})_n) \geq 1$.

5 Hilbert series of the Hopf algebra WMat

In this section, we compute the number of packed words with length n and supremum k . It is the same as the number of ordered partitions of an n -element set. Using the formula of Stirling numbers of the second kind, one can get the explicit formula for the number of packed words with length n , number which we denote by d_n .

Definition 5.1 *The Stirling numbers of the second kind count the number of set partitions of an n -element set into precisely k parts. The Stirling numbers, denoted by $S(n, k)$ are given by the recursive definition:*

1. $S(n, n) = 1 (n \geq 0)$,
2. $S(n, 0) = 0 (n > 0)$,

$$3. S(n+1, k) = S(n, k-1) + kS(n, k), \text{ for } 0 < k \leq n.$$

One can define a word by its positions, that means if a word $w = x_{i_1}x_{i_2}\dots x_{i_n}$ has length n and $IAlph(w) = \{1, 2, \dots, k\}$, then this word can be determine from the list $[S_1, S_2, \dots, S_k]$, where the S_i 's is a partition of $\{1, 2, \dots, n\}$, and S_i is the set of positions of x_i in the word w , with $1 \leq i \leq k$.

One can divide the set of packed words with length n and supremum k in two part: "pure" packed words (which have no x_0 in their alphabet), denote $pack_{n,k}^+(X)$ and packed words which have x_0 in their alphabet, denote $pack_{n,k}^0(X)$. It is clear that:

$$d(n, k) = \#pack_{n,k}^+(X) + \#pack_{n,k}^0(X). \quad (44)$$

Let us now compute the cardinal of these two sets $pack_{n,k}^+(X)$ and $pack_{n,k}^0(X)$.

Consider a word $w \in pack_{n,k}^+(X)$, then $IAlph(w) = \{1, 2, \dots, k\}$. This word is determined by $[S_1, S_2, \dots, S_k]$, in which S_i is a set of positions of x_i , for $1 \leq i \leq k$. One can see that:

1. $S_i \neq \emptyset$,
2. $\sqcup_{1 \leq i \leq k} S_i = \{1, 2, \dots, n\}$.

Thus, one has the cardinal of packed words with length n and supremum k :

$$d^+(n, k) = \#pack_{n,k}^+(X) = S(n, k)k!. \quad (45)$$

Similarly, a word $w \in \#pack_{n,k}^0(X)$ can be determined by $[S_0, S_1, S_2, \dots, S_k]$ where S_i is the set of positions of x_i , for all $0 \leq i \leq k$. One then has:

$$d^0(n, k) = \#pack_{n,k}^0(X) = S(n, k+1)(k+1)!. \quad (46)$$

From the two equations above, one can get the number of packed word with length n , supremum k :

$$d(n, k) = d^+(n, k) + d^0(n, k) = S(n, k)k! + S(n, k+1)(k+1)!. \quad (47)$$

From this formula, using Maple, one can get some values of $d(n, k)$. We give in the following table the first values.

		k								
		0	1	2	3	4	5	6	7	8
n	0	1	0	0	0	0	0	0	0	0
	1	1	1	0	0	0	0	0	0	0
	2	1	3	2	0	0	0	0	0	0
	3	1	7	12	6	0	0	0	0	0
	4	1	15	50	60	24	0	0	0	0
	5	1	31	180	390	360	120	0	0	0
	6	1	63	602	2100	3360	2520	720	0	0
	7	1	127	1932	10206	25200	31920	20160	5040	0
	8	1	255	6050	46620	166824	317520	332640	181440	40320

Table 1: Values of $d(n, k)$ given by the explicit formula (47) and computed with Maple.

The formula of d_n , $n \geq 1$ is then given by

$$\begin{aligned}
d_n &= \sum_{k=0}^n d(n, k) = \sum_{k=0}^n [S(n, k)k! + S(n, k+1)(k+1)!] = \sum_{k=0}^n S(n, k)k! + \sum_{k=0}^n S(n, k+1)(k+1)! \\
&= \begin{cases} 1 & \text{if } n = 0, \\ 2 \sum_{k=1}^n S(n, k)k! & \text{if } n \geq 1 \end{cases} \quad (\text{note that } S(n, 0) = S(n, n+1) = 0). \quad (48)
\end{aligned}$$

Using again Maple, one can get the following table:

n	0	1	2	3	4	5	6	7	8	9	10
d_n	1	2	6	26	150	1082	9366	94586	1091670	14174522	204495126

Table 2: Some value of d_n by the formula (48).

Recall that one has:

$$e^{y(e^x-1)} = \sum_{k=0}^{\infty} \sum_{n=k}^{\infty} S(n, k) \frac{x^n y^k}{n!} .. \quad (49)$$

From equations (45) and (49), one has:

$$e^{y(e^x-1)} = \sum_{k=0}^{\infty} \sum_{n=k}^{\infty} d^+(n, k) \frac{x^n y^k}{n! k!} . \quad (50)$$

Partially derivating equation (49) with respect to y, one has:

$$(e^x - 1)e^{y(e^x-1)} = \sum_{k=1}^{\infty} \sum_{n=k}^{\infty} S(n, k) k \frac{x^n y^{k-1}}{n!} . \quad (51)$$

From equations (46) and (51), one has:

$$(e^x - 1)e^{y(e^x-1)} = \sum_{k=1}^{\infty} \sum_{n=k+1}^{\infty} d^0(n, k) \frac{x^n y^k}{n! k!} . \quad (52)$$

One can thus find the number of packed words with length n and supremum k as the coefficient of degree (n,s) in the Taylor series of the function e^{ye^x+x-y} .

We have defined here a new non-commutative and non-cocommutative bialgebra on certain words, based on the selection/quotient principle. The next step would be to find a polynomial realization of it.

Acknowledgements

The quality of this paper was greatly enhanced by the gracious assistance of Prof. Gérard H.E. Duchamp and Adrian Tanasa, reading teachers, who have taught me most of what I know about this. I would also like to thank Prof. Jean-Yves Thibon, Gaspard-Monge Institute at University Marne-la-Vallée , for various discussions and suggestions.

References

- [1] A. CONNES, D. KREIMER, *Renormalization in quantum field theory and the Riemann-Hilbert problem I: The Hopf algebra structure of graphs and the main theorem*, Commun. Math. Phys. **210** (2000), no. 1, 249-273.
- [2] G.H.E. DUCHAMP, F. HIVERT, J.Y. THIBON : *Non commutative functions VI: Free quasi-symmetric functions and related algebras* IJAC Vol. 12 **5** (2002).
- [3] I.M. GELFAND, D. KROB, A. LASCoux, B. LECLERC, V.S. RETAKH, J.Y. THIBON, *Noncommutative symmetric functions*, Adv. Math. **112** (1995), 218-348.
- [4] J.-L. LODAY, M. O. RONCO, *Trialgebras and families of polytopes*, Homotopy Theory: Relations with Algebraic Geometry, Group Cohomology, and Algebraic K-theory, Contemporary Mathematics 346, (2004), AMS.
- [5] J.- L. LODAY, M. O. RONCO, *Hopf algebra of the Planar Binary Trees*, Adv. Math. **139** (1998), 293-309.

- [6] C. MALVENUTO, C. REUTENAUER, *Duality between Quasi-Symmetric Functions and the Solomon Descent Algebra*, Journal of Algebra **177**, 967-982 (1995).
- [7] F. MARKOPOULOU, *Coarse graining in spin foam models*, Class. Quant. Grav. **20** (2003) 777-800 [gr-qc/0203036].
- [8] A. TANASA, D. KREIMER, *Combinatorial Dyson-Schwinger equations in noncommutative field theory*, J. Noncomm. Geom. (in press) arXiv:0907.2182 [hep-th].
- [9] A. TANASA, F. VIGNES-TOURNERET, *Hopf algebra of non-commutative field theory*, J. Noncomm. Geo. Volume 2, Issue 1, 2008, 125 - 139.
- [10] A. TANASA, *Combinatorial Hopf Algebras in (Noncommutative) Quantum Field Theory*, Rom. J. Phys. **55**, 1142 (2010) [arXiv:1008.1471 [math.CO]].
- [11] A. TANASA, *Algebraic structures in quantum gravity*, Class. Quant. Grav. **27**, 095008 (2010) [arXiv:0909.5631 [gr-qc]].
- [12] W. SCHMITT, *Incidence Hopf algebras*, Journal of Pure and Applied Algebra **96** (1994), 299-330.
- [13] H. WHITNEY, *On the abstract properties of linear dependence*, Amer. J. Math. **57** (1935), 509-533.